

Analysing the complexity of marine food webs using amino acid stable nitrogen isotopes: from trophic positions to diets

A. Bode (February 2022)

References

- Basedow SL, de Silva NAL, Bode A, van Beusekorn J (2016) Trophic positions of mesozooplankton across the North Atlantic: estimates derived from biovolume spectrum theories and stable isotope analyses. *J Plankton Res* 38:1364-1378 (example of microbial vs. metazoan food web)
- Bradley CJ, Wallsgrove NJ, Choy CA, Drazen JC, Hetherington ED, Hoen DK, Popp BN (2015) Trophic position estimates of marine teleosts using amino acid compound specific isotopic analysis. *Limnol Oceanogr Methods* 13:476-493 (review of TDF and TP in fish)
- Bode A, Besada V, Pérez-Fernández B, Viñas L (2021d) Amino acid $\delta^{15}\text{N}$ can detect diet effects on pollution risks for Yellow-Legged Gulls overlooked by trophic position. *Frontiers in Marine Science* 8: 657431 <https://doi.org/10.3389/fmars.2021.657131> (example of diet tracers using CSIA-AA)
- Bode A, Olivar MP, Hernández-León S (2021a) Trophic indices for micronektonic fishes reveal their dependence on the microbial system in the North Atlantic. *Scientific Reports* 11:8488 <https://doi.org/10.1038/s41598-41021-87767-x> (example of microbial contributions to TP)
- Bode A, Olivar MP, López-Pérez C, Hernández-León S (2021b) The microbial contribution to the trophic position of stomiiform fishes. *ICES J Mar Sci* 78:3245-3253 <https://doi.org/10.1093/icesjms/fsab189> (example of microbial contributions to TP)
- Bode A, Olivar MP, López-Pérez C, Hernández-León S (2021c) Stable nitrogen isotopes reveal microbial contribution to the trophic position of micronektonic fishes. *ASLO 2021. Association for the Sciences of Limnology and Oceanography, web conference, 22-27 June 2021* (example of microbial contributions to TP)
- Cherel Y, Bustamante P, Richard P (2019) Amino acid $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ from sclerotized beaks: a new tool to investigate the foraging ecology of cephalopods, including giant and colossal squids. *Mar Ecol Prog Ser* 624:89-102 (example of application of C CSIA-AA)
- Chikaraishi Y, Ogawa NO, Kashiyama Y, Takano Y, Suga H, Tomitani A, Miyashita H, Kitazato H, Ohkouchi N (2009) Determination of aquatic food-web structure based on compound-specific nitrogen isotopic composition of amino acids. *Limnol Oceanogr Methods* 7:740-750 (reference method for TP estimations from CSIA-AA)
- Decima M, Landry MR, Bradley CJ, Fogel ML (2017) Alanine $\delta^{15}\text{N}$ trophic fractionation in heterotrophic protists. *Limnol Oceanogr* 62:2308-2322 (basis for determination of microbial trophic steps using CSIA-AA)
- Decima M, Landry M (2020) Resilience of plankton trophic structure to an eddy-stimulated diatom bloom in the North Pacific Subtropical Gyre. *Mar Ecol Prog Ser* 643:33-48 (example of microbial vs. metazoan contributions to TP)
- Gutiérrez-Rodríguez A, Décima M, Popp BN, Landry MR (2014) Isotopic invisibility of protozoan trophic steps in marine food webs. *Limnol Oceanogr* 59:1590-1598 (experimental demonstration of CSIA-AA tracers for microbial food webs)

- Hannides CCS, Popp BN, Choy CA, Drazen JC (2013) Midwater zooplankton and suspended particle dynamics in the North Pacific Subtropical Gyre: A stable isotope perspective. *Limnol Oceanogr* 58:1931-1946 (example of application of C and N CSIA-AA)
- Ishikawa NF, Chikaraishi Y, Takano Y, Sasaki Y, Takizawa Y, Tsuchiya M, Tayasu I, Nagata T, Ohkouchi N (2018) A new analytical method for determination of the nitrogen isotopic composition of methionine: Its application to aquatic ecosystems with mixed resources. *Limnol Oceanogr Methods* 16:607-620 (use of CSIA-Met as tracer of source primary producers)
- Larsen T, Hansen T, Dierking J (2020) Characterizing niche differentiation among marine consumers with amino acid $\delta^{13}\text{C}$ fingerprinting. *Ecol Evol* 10:7768-7782 (application of C CSIAA)
- McCarthy MD, Benner R, Lee C, Fogel ML (2007) Amino acid nitrogen isotopic fractionation patterns as indicators of heterotrophy in plankton, particulate, and dissolved organic matter. *Geochim Cosmochim Acta* 71:4727-4744 (definition and use of microbial heterotrophy degradation indices from CSIA-AA)
- McCarthy MD, Lehman J, Kudela R (2013) Compound-specific amino acid $\delta^{15}\text{N}$ patterns in marine algae: Tracer potential for cyanobacterial vs. eukaryotic organic nitrogen sources in the ocean. *Geochim Cosmochim Acta* 103:104-120 (one reference method for derivatization for CSIA-AA)
- McClelland JW, Montoya JP (2002) Trophic relationships and the nitrogen isotopic composition of amino acids in plankton. *Ecology* 83:2173-2180 (first application of CSIA-AA to marine plankton, early definition of trophic and source AAs)
- McMahon KW, McCarthy MD (2016) Embracing variability in amino acid $\delta^{15}\text{N}$ fractionation: mechanisms, implications, and applications for trophic ecology. *Ecosphere* 7:e01511, <https://doi.org/01510.01002/ecs01512.01511> (review of TDF for N in CSIA-AA, including multi-TDF)
- Mompeán C, Bode A, Gier E, McCarthy MD (2016) Bulk vs. aminoacid stable N isotope estimations of metabolic status and contributions of nitrogen fixation to size-fractionated zooplankton biomass in the subtropical N Atlantic. *Deep Sea Res* 114:137-148 (example of application of CSIA-AA to trace diazotrophy)
- Nielsen JM, Popp BN, Winder M (2015) Meta-analysis of amino acid stable nitrogen isotope ratios for estimating trophic position in marine organisms. *Oecologia* 178:631-642 (review of TDF for CSIA-AA)
- Ohkouchi N, Chikaraishi Y, Close HG, Fry B, Larsen T, Madigan DJ, McCarthy MD, McMahon KW, Nagata T, Naito YI, Ogawa NO, Popp BN, Steffan S, Takano Y, Tayasu I, Wyatt ASJ, Yamaguchi YT, Yokoyama Y (2017) Advances in the application of amino acid nitrogen isotopic analysis in ecological and biogeochemical studies. *Org Geochem* 113:150-174 (review of recent developments of TP estimations from CSIA-AA and error propagations)
- Popp BN, Graham BS, Olson RJ, Hannides CCS, Lott MJ, López-Ibarra GA, Galván Magaña F, Fry B (2007) Insight into the trophic ecology of Yellowfin Tuna, *Thunnus albacares*, from compound-specific nitrogen isotope analysis of proteinaceous amino acids. In: Dawson TE, Siegwolf RTW (eds) *Stable isotopes as indicators of ecological change*, Vol 1. Elsevier, p 173-190 (early definition of trophic and source AAs, application to TP estimations of top marine predator)

- Ramirez MD, Besser AC, Newsome SD, McMahon KW (2021) Meta-analysis of primary producer amino acid $\delta^{15}\text{N}$ values and their influence on trophic position estimation. *Methods Ecol Evol* <https://doi.org/10.1111/2041-210X.13678> (review of CSIA-AA beta for primary producers, use of beta-mix)
- Ruiz-Cooley RI, Gerrodette T, Fiedler PC, Chivers SJ, Danil K, Ballance LT (2017) Temporal variation in pelagic food chain length in response to environmental change. *Sci Adv* 3: e1701140 <https://doi.org/10.1126/sciadv.1701140> (example of change in TP, FCL, using CSIA-AA)
- Shen Y, Guilderson TP, Sherwood OA, Castro CG, Chavez FP, McCarthy MD (2021) Amino acid $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ patterns from sediment trap time series and deep-sea corals: Implications for biogeochemical and ecological reconstructions in paleoarchives. *Geochim Cosmochim Acta* 297:288-307 (example of use of C and N CSIA-AA)
- Takizawa Y, Takano Y, Choi B, Dharampal PS, Steffan SA, Ogawa NO, Ohkouchi N, Chikaraishi Y (2020) A new insight into isotopic fractionation associated with decarboxylation in organisms: implications for amino acid isotope approaches in biogeoscience. *Prog Earth Planet Sci* 7:50 <https://doi.org/10.1186/s40645-020-00364-w> (comparison of C and N measurements using CSIA-AA)
- Twining CW, Taipale SJ, Ruess L, Bec A, Martin-Creuzburg D, Kainz MJ (2020) Stable isotopes of fatty acids: current and future perspectives for advancing trophic ecology. *Philos Trans R Soc B-Biol Sci* 375:1804 <https://doi.org/10.1098/rstb.2019.0641> (review of CSIA-FA methods and applications)
- Vander Zanden MJ, Fetzer W (2007) Global patterns of aquatic food chain length. *Oikos* 116:1378-1388 (applications and ecological implications of FCL)
- Viñas, L., Besada, V., Pérez-Fernández, B., Bode, A., 2020. Yellow-legged gull eggs (*Larus michahellis*) as persistent organic pollutants and trace metal bioindicator for two nearby areas with different human impact. *Environ Res* 190(110026), <https://doi.org/10.1016/j.envres.2020.110026> (example of problem requiring CSIA-AA)
- Yun HY, Lee D-H, Choi H, Won E, Shin K-H (2020) Evaluation of the quantity and the carbon isotopic composition of amino acids by using diverse sample residues after lipid extraction. *Anal Bioanal Chem* 412:4383-4391 (review of CSIA-FA methods)
- Xing D, Choi B, Takizawa Y, Fan R, Sugaya S, Tuchiya M, Ohkouchi N, Chikaraishi Y (2020) Trophic hierarchy of coastal marine fish communities viewed via compound-specific isotopic analysis of amino acids. *Mar Ecol Prog Ser* 652:137-144 (example of baseline identification using CSIA-AA)