Increase in membrane thickness during development compensates for eggshell thinning due to calcium uptake by the embryo in falcons

Aurora M. Castilla & Stefan Van Dongen & Anthony Herrel & Amadeu Francesch & Juan Martínez de Aragón & Jim Malone & Juan José Negro

Abstract We compared membrane thickness of fully developed eggs with those of non-developed eggs in different endangered falcon taxa. To our knowledge, membrane thickness variation during development has never been examined before in falcons or any other wild bird. Yet, the egg membrane constitutes an important protective barrier for the developing embryo. Because eggshell thinning is a general process that occurs during bird development, caused by calcium uptake by the embryo, eggs are expected to be less protected and vulnerable to breakage near the end of development. Thus, egg membranes could play an important protective role in the later stages of development by getting relatively thicker.

A. M. Castilla (★)
Estación Biológica y de Cultura Científica de Sanaüja,
Ap. Correos nº 35,
25280 Solsona, Lleida, Spain
e-mail: castilla.aurora@gmail.com
e-mail: castilla@ebd.csic.es

A. M. Castilla
Estación Biológica de Doñana, Agencia Estatal Consejo Superior de Investigaciones Científicas (CSIC),
C/Americo Vespucio s/n,
41092 Sevilla, Spain

S. Van Dongen Département d'Ecologie et de Gestion de la Biodiversité, UMR 7179 C.N.R.S/M.N.H.N, 57 rue Cuvier, Case postale 55, 75231 Paris Cedex 5, France e-mail: stefan.vandongen@ua.ac.be

A. Herrel

Evolutionary Biology Group, Department of Biology, University of Antwerp, Groenenborgerlaan 171, 2020 Antwerp, Belgium e-mail: anthony.herrel@mnhn.fr We used linear mixed models to explore the variation in membrane thickness (n=378 eggs) in relation to developmental stage, taxon, female age, mass and identity (73 females), egg-laying sequence (105 clutches) and the study zone. Our results are consistent with the prediction that egg membranes are thicker in fully developed eggs than in nondeveloped eggs, suggesting that the increase in membrane thickness during development may compensate for eggshell thinning. In addition, our data shown that thicker membranes are associated with larger, heavier and relatively wider eggs, as well as with eggs that had thinner eggshells. Egg-laying sequence, female age and the study zone did not explain the observed variation of membrane thickness in

A. Francesch

Unitat de Genètica Avícola, Centre de Mas Bové; IRTA, Apartat 415, 43280 Reus, Tarragona, Spain e-mail: amadeu.francesch@irta.es

J. Martínez de Aragón Centre Tecnològic Forestal de Catalunya, Carretera de Sant Llorenç de Morunys Km 2, 25280 Solsona, Lleida, Spain e-mail: mtzda@ctfc.es

J. Malone

Information Services Librarian, Weill Cornell Medical College in Qatar, P.O. Box 24144, Doha, Qatar e-mail: jrm2006@qatar-med.cornell.edu

J. José Negro

Department of Evolutionary Ecology, Estación Biológica de Doñana, Agencia Estatal Consejo Superior de Investigaciones Científicas, Americo Vespucio s/n, 41092 Sevilla, Spain e-mail: negro@ebd.csic.es the falcon taxa studied. As we provide quantitative data on membrane thickness variation during development in falcons not subjected to contamination or food limitation (i.e. bred under captive conditions), our data may be used as a reference for studies on eggs from natural populations. Considering the large variation in membrane thickness and the multiple factors affecting on it and its importance in the protection of the embryo, we encourage other researchers to include measurements on membranes in studies exploring eggshell thickness variation.

Keywords Egg membrane · Eggshell thickness · Egg-laying sequence · Hybrid · Conservation · Raptor · CITES

Introduction

The avian egg provides the developing embryo with an ideal environment for development (Romanoff and Romanoff 1949; Rahn and Ar 1974; Solomon 1991). The eggshell membrane is a smooth layer of homogeneous dense material that functions as a site for respiration, nutrient transport, and waste storage during embryonic development (Kutchai and Steen 1971; Peakall et al. 1983; Cobb et al. 2003; Pepper et al. 2004). Membranes also prevent the embryo from drying out (Mao et al. 2007) and facilitate transport of calcium during late embryonic stages (Elaroussi and Deluca 1994; Ecay et al. 2004). Membranes also act as a filter against the penetration of microorganisms into the egg (Ahlborn and Sheldon 2005; Chao et al. 2007).

Membrane thickness has received very little attention in the literature (but see Yoshizaki and Saito 2002; Mao et al. 2007; Feberwee et al. 2009) when compared to eggshell thickness. Eggshell thinning has been the focus of much research in falcons, mainly because of the reproductive failure that have occurred in many parts of the northern hemisphere from about 1950 possibly due to persistent pollutants (Cade et al. 1971; Bunck et al. 1985; Bennett 1995; Peakall and Lincer 1996; Wegner et al. 2005; Falk et al. 2006; Kamata et al. 2006; Jagannath et al. 2008; Castilla et al. 2009a). Embryo development, however, also causes eggshell thinning in many bird species (Vanderstoep and Richards 1970; Bunck et al. 1985; Castilla et al. 2007; Karlsson and Clas Lilja 2008; Fernie et al. 2009) including falcons of different taxa (Castilla et al. 2010), as the embryos utilise minerals and calcium from the eggshell to assist in the development of the skeleton, muscles and brain (Tuan et al. 1991; Blom and Lilja 2004; Wilkin et al. 2009). One means to compensate for eggshell thinning in some birds is the deposition of a thicker layer of shell pigment (Gosler et al. 2005). Another way to compensate for eggshell thinning that would provide protection for the

embryo during the latest phase of their development would be to increase the thickness of the egg membranes.

In this study, we examined the effect of developmental stage on membrane thickness variation in different falcon taxa. We tested the prediction that membranes from fully developed eggs are thicker than those from non-developed eggs. We additionally explored the effect of different egg characteristics, female traits, egg-laying sequence and study zone on membrane thickness variation.

Materials and methods

Study animals and eggs

Falcons are included in the list of the Convention of International Trade of Endangered Species (CITES; Tucker and Heath 1994). In addition, gaining access to their nests and eggs is very difficult because they are normally in sharp cliffs. Consequently, there is a large gap in our knowledge regarding membrane or eggshell thickness variation and the factors affecting on it.

The falcon species examined in this study are protected, rare or endangered and include the peregrine falcon (Falco peregrinus peregrinus; P), the red shaheen falcon (Falco peregrinus babylonicus; R); the intraspecific hybrid peregrine \times red shaheen (F. peregrinus peregrinus \times F. peregrinus babylonicus; PR); the saker falcon (Falco cherrug; S), the gyr falcon (Falco rusticolus), and the interspecific hybrid between the saker falcon (F. cherrug) and the gyr falcon (F. rusticolus; SG). Females show large differences in body size (ranging from 600 to 1,800 g) depending on the taxon (Cramp and Simmons 1980). In this study, body mass was measured at the end of summer after reproduction was finished. The birds were captured in their cages and weighed with an electronic balance to the nearest gramme (KRUPS 840, Germany). The age of the females was provided by the one responsible for the breeding centre "Roc Falcon" who has information on the birth date of all individuals.

The eggs were obtained from two different breeding zones in Catalonia (NE Spain) that were not subjected to agricultural or industrial pollution. One zone was near the coast (GI) at 97 m.a.s.l. (mean annual temperature=14°C, mean annual precipitation=650 mm, mean relative humidity=80%); the other was in the Pyrenees mountains (RF) at 800 m.a.s.l. (mean annual temperature=12°C, mean annual precipitation= 650 mm, mean relative humidity=65%).

Falcons are bred worldwide and a common target for conservation programmes (Rahbek 1993). This provides a unique opportunity where a large sample of eggs can be obtained for scientific research. Moreover, falcons in captive breeding facilities are bred under optimal conditions and fed high quality food. Since all taxa and individuals are treated similarly, results can be compared without bias in that respect. We examined 105 clutches from 73 different females (total of 211 non-developed and 167 developed eggs) that were collected between March and July of 2007.

In the field, the falcon species studied here typically have one or two clutches from four to six eggs and an incubation period that last ca. 25 days (Cramp and Simmons 1980). Egg pulling (i.e. removing eggs as they are laid) was conducted in both zones, so females did not produce true clutches as they would do in the wild. However, most females produced between four and nine eggs. Data on egglaying sequence was obtained by writing down a number on the eggshell as the female was laying it. The eggshells from hatched chicks were obtained from the birth rooms.

We used an ovoscope and direct observations after egg breakage to establish two developmental categories: eggs with non-developed embryos (i.e. infertile or aborted during the first week of incubation; Fig. 1), and developed eggs (i.e. hatched eggs or aborted eggs with fully developed embryos). Egg size (length and width) was measured with a calliper (Mitutoyo) to the nearest 0.01 mm. Egg mass was measured in fresh eggs that were recently laid with an electronic Sartorious AG, balance Goettingen, Germany (to 0.01 g). Membranes were separated from the eggshells after immersing them in a plastic box with water for 10 min. In eggs of both developmental stages, we only measured the outer thick eggshell membrane after cleaning it with water and removing the thin vascularized chorioallantoic membrane (Fig. 1). Membrane thickness was measured using a micrometer (Mitutoyo) to the nearest 0.001 mm. Measurements were taken in a similar way, that is, by compressing the membrane with a standardised pressure provided by the spring in the micrometer. Membrane and eggshell thickness were measured in one site around the equator of the egg. All measurements were done by the same person; we are confident that these are perfectly comparable among species.

Statistical procedures

Because all egg dimensions (length, width, mass and shell thickness) as well as female mass were highly correlated (r > 0.4 and p < 0.0001 in all cases), we based all analyses on a multivariate summary of the these data. We first performed a principal component analysis to reduce the dimensionality of the data set. This resulted in a new set of uncorrelated variables that can be analysed separately. Results of the principal component analysis were represented graphically using biplots, where the cosine of the angle between two vectors provides an estimate of the correlation between the respective variables. Principal components were used as independent variables in a mixed



Fig. 1 Eggshells, membranes and embryos of different developmental stages in falcon species. a, d, e Peregrine falcon (Falco peregrinus peregrinus). b Intraspecific hybrid peregrine×red shaheen (F. peregrinus peregrinus×F. peregrinus babylonicus). c Red shaheen falcon (Falco peregrinus babylonicus). f Interspecific

hybrid saker×gyr (Falco cherrug×Falco rusticolus). The chorioallantoic membrane (vascularized) is present in nearly fully developed embryos (b, c) and in hatched eggs (d, right). Membranes are also shown in infertile eggs (d left, e, f)

linear model with membrane thickness as dependent variable. Additionally, developmental stage and zone were added as fixed factors. Laying sequence and female age were added as continuous covariates. Female identity was added as a random effect to estimate among-female variation and to take the dependency of the data for eggs from a single female into account. Tests of fixed effects were based on F tests with degrees of freedom approximated by Kenward and Rogers method. The random female effect was tested using a likelihood ratio test. The data was analysed using linear mixed models in SAS (version 9.2).

In this study, we have compared individuals of different species that cannot be treated as statistically independent data points following the premise put forth by Joe Felsenstein (1985, 2008). However, the number of falcon species that we studied here is too low (n=3), thus providing very few degrees of freedom if one were to do an independent contrast analysis on the species means. Moreover, some of our data points are hybrids, making it conceptually hard to include them as if they were 'independent' species. We have to be aware of this limitation in the present study.

Results

The smaller falcons of the peregrine group (P, R, PR) have egg membranes significantly thinner than the big falcons (S, G, SG) when considering both developmental stages separately (developed: analysis of variance (ANOVA), $F_{1,208} = 26.76$, p<0.001; undeveloped: ANOVA, $F_{1,166} =$ 9.16, p=0.003; Table 1, Fig. 2). We also found a high significant correlation between membrane thickness and eggshell thickness in eggs under different developmental stages: undeveloped eggs (regression on Log₁₀-transformed data: r=0.37; p<0.001; slope=0.94; intercept=-0.71) and fully developed eggs (regression on Log₁₀-transformed data: r=0.32; P<0.001; slope=0.60; intercept=-0.79; Fig. 2).

Because several of the morphometric measurements were highly correlated, we conducted first a principal component analysis, and found that the first three components explained 93% of all variation (Table 2). Eigenvectors and a graphical representation of this principal component analysis as biplots are given in Table 2 and Fig. 3. The first principal component corresponded to an overall size measure (egg and female size). The second principal component was determined by a contrast between egg length and shell thickness. Higher scores for the second component correspond to relative long eggs with relative thin shell. Finally, the third principal component was determined by a contrast between egg length and shell thickness on the one hand and egg width and female mass on the other hand. The strong correlations showed by the principal component analysis allow imputing the missing data by using multiple imputations. After that, all data can be used to see which factors are correlated with membrane thickness (Table 3).

Table 1 Measurements of membrane thickness (in millimetres) for eggs with fully developed embryos or with non-developed embryos in different falcon taxa

Taxa	Zone	Developed eggs					Non-developed eggs				
		Mean	SD	Max	Min	Ν	Mean	SD	Max	Min	Ν
Р	GI	0.088	0.006	0.094	0.077	9	0.066	0.015	0.1	0.037	32
	RF	0.077	0.016	0.101	0.036	29	0.059	0.019	0.095	0.019	40
R	GI	0.071	0.015	0.092	0.035	14	0.054	0.018	0.09	0.025	28
	RF	0.061	0.018	0.081	0.026	7	0.051	0.020	0.087	0.02	16
PR	GI	0.066	0.019	0.105	0.038	10	0.059	0.009	0.065	0.044	6
	RF	_	_	_	_						
S	GI	0.070	0.011	0.093	0.054	9	0.069	0.014	0.098	0.048	15
	RF	0.083	0.014	0.102	0.053	30	0.074	0.012	0.094	0.057	20
G	GI	_	_	_	_						
	RF	_	_	_	_		0.076	0.007	0.088	0.066	9
SG	GI	0.072	0.013	0.091	0.054	6	0.081	0.007	0.091	0.073	7
	RF	0.084	0.013	0.106	0.054	53	0.068	0.020	0.108	0.032	38

Indicated are the means and standard deviations (SD), the maximum and minimum values, the sample size (N) and the zone (see "Materials and methods")

P peregrine falcon (Falco peregrinus peregrinus, R=red shaheen falcon (Falco peregrinus babylonicus), PR intraspecific hybrid peregrine×red shaheen (F. peregrinus peregrinus ×F. peregrinus babylonicus), S saker falcon (Falco cherrug), G gyr falcon (Falco rusticolus), SG interspecific hybrid saker×gyr (F. cherrug×F. rusticolus)



Fig. 2 Membrane thickness (mean±SE) of eggs of different developmental stage (white=undeveloped; black=fully developed) for the different falcons. P peregrine falcon (Falco peregrinus peregrinus), R red shaheen falcon (Falco peregrinus babylonicus), PR intraspecific hybrid peregrine×red shaheen (F. peregrinus peregrinus \times F. peregrinus babylonicus), S saker falcon (Falco cherrug), G gyr falcon (Falco rusticolus), SG interspecific hybrid saker×gyr (F. cherrug×F. rusticolus)

Membranes were significantly thicker in eggs at a later developmental stage where fully developed eggs had thicker membranes than non-developed eggs (Fig. 2, Table 3). For all falcon taxa, membrane thickness was also correlated with egg size and female body mass (i.e. a positive correlation with the first principal component, Figs. 3 and 4), and this correlation did not differ between the two developmental stages (i.e. no two-way interaction, Table 3). Thicker membranes corresponded to relatively long eggs with thinner eggshells (i.e. correlation with PC2, Table 3, Fig. 4). Species, zone, egg-laying sequence and the third principal component did not explain the observed membrane thickness variation in falcon taxa (Table 3, Fig. 5).

Discussion

In this study, we provide the first quantitative data on membrane thickness variation during development for non-



Fig. 3 Relationship between eggshell thickness and membrane thickness for undeveloped eggs (a) and for fully developed eggs (b). Black circle: peregrine falcon, P (Falco peregrinus peregrinus); white square: red shaheen falcon, R (Falco peregrinus babylonicus); red triangle up: intraspecific hybrid peregrine × red shaheen, PR (F. peregrinus peregrinus × F. peregrinus babylonicus); green triangle down: saker falcon, S (Falco cherrug); yellow diamond: gyr falcon, G (Falco rusticolus), blue hexagon: interspecific hybrid saker × gyr, SG (F. cherrug × F. rusticolus)

poultry birds. However, as the falcons in our study were raised in optimal conditions, this allows our data to be used as a reference for studies on eggs from natural populations. Our results are consistent with the prediction that membrane thickness increases during incubation and suggest an

Table 2 Eigenvectors and proportion variance explained by the principal component analysis of egg characteristics and female mass (a screen plot is given in Fig. 3)

	PC1	PC2	PC3	PC4	PC5
Length	0.400928	0.764996	0.337077	0.104549	0.359843
Width	0.463732	-0.155637	-0.390096	-0.682461	0.377892
Egg mass	0.507505	0.175252	-0.077946	-0.111582	-0.832586
Shell thickness	0.404397	-0.559875	0.720487	0.022831	0.058141
Female mass	0.450679	-0.215372	-0.457195	0.714384	0.17644
% Variance explained	70	12	9	6	3
Cumulative %	70	82	91	97	100

Table 3 Overview of significance tests of associations between m embrane thickness in different falcon taxa and explanatory variables

Parameter	F value	df 1	df 2	p value
PC1	30.0	1	89	< 0.0001
PC2	10.4	1	302	0.001
PC3	1.99	1	354	0.16
Developmental stage	67.8	1	365	< 0.0001
Egg-laying sequence	0.01	1	53	0.90
Female age	0.44	1	59	0.51
Zone	1.70	1	54	0.20
PC1×developmental stage	1.57	1	349	0.21
PC2×developmental stage	0.05	1	353	0.82
PC3×developmental stage	0.09	1	336	0.76

important effect on the protection of the developing embryo. This should be especially relevant for bird species nesting in rocky areas, such as falcons (Cramp and Simmons 1980). The egg membrane can act to prevent global cracking of the eggs in case of small cracks, as it provides a strong coherence among the parts of the eggshell. A small crack in the eggshell need not be fatal to the embryo. As long as the egg membrane is intact, the embryo is shielded from external influences such as pathogens and some fluctuation in environmental conditions. A similar trend towards an increase in membrane thickness during development was observed in the redlegged partridge (Alectoris rufa; Castilla and Herrel, unpublished). Our results are also in agreement with the findings of a recent study (Feberwee et al. 2009) which revealed that when the mammillary layer of the calcified zone of the egg was absent, the inner eggshell membranes were thicker.

In contrast to our observations, shell membranes in farm quails become thinner during incubation (Yoshizaki and Saito 2002), which has been suggested to be related to the need to satisfy the requirements for gas exchange (Kutchai and Steen 1971). However, to make comparisons across studies and species, similar methods should be employed since the structure of egg membranes (as demonstrated by electron micrographs) is different in the region of the air chamber compared to other regions of the egg (Mao et al. 2007). In our study, we used the same micrometer, and we measured membrane thickness always in the same site around the equator in all eggs. Also, the measurements of developed and non-developed eggs were conducted during the same time. Thus, our results showing the same trend of membrane thickness increase during development in all the five falcon taxa examined are not likely the result of a measurement artefact.



Fig. 4 Biplots of the three first principal components of a principal component analysis of egg measurements and female mass. Data from the different

species are indicated by different

numbers: 1=peregrine falcon

(Falco peregrinus peregrinus);

2=red shaheen falcon (Falco

peregrinus babylonicus); 3=

intraspecific hybrid peregrine× red shaheen (F. peregrinus peregrinus×F. peregrinus babylonicus); 4=saker falcon (Falco cherrug); 5=gyr falcon (Falco rusticolus); 6= interspecific hybrid saker×gyr (F. cherrug×F. rusticolus)

D Springer







Fig. 5 Associations between membrane thickness and the first and second principal component of egg measurements and female mass for fully developed (grey) and non-developed (black) eggs in different falcon taxa. Different species are indicated by different numbers: 1=peregrine falcon (Falco peregrinus peregrinus); 2=red

In contrast to our expectations, egg-laying sequence did not have an effect on membrane thickness. Yet, a decrease in mean shell thickness in later clutches of captive peregrine falcons (Burnham et al. 1984; Falk and Møller 1990) and in other falcon taxa (Castilla et al. 2010) has been observed. An additional factor that could influence membrane thickness variation with respect to egg-laying sequence is variation in chick mass related to sex (females are much bigger in falcons). However, since information on the sex of the developing embryos was not available, this needs to be investigated further.

We found that heavier females produced larger and heavier eggs with thicker membranes. Our results are in agreement with other authors who hypothesised that for an effective incubation without breaking the eggs in the nests, big birds have to produce bigger and thicker eggs that small birds (Ar et al. 1979). We also found that the age of the female did not influence on membrane thickness variation in this study. This result was rather surprising since a previous study on falcons (albeit using infertile non-developed eggs) found that older females produced eggs with thinner membranes than younger females (Castilla et al. 2009b). Because we used different females of different age and condition in two studies, this may have influenced the different trend in our results with younger females being included in the present study. However, this variation is biologically relevant and important and should be taken into consideration. Additional data on how female age influences egg and egg membrane characteristics are needed to gain a fuller understanding of the observed patterns.

In our study, membrane thickness variation among taxa

hybrid peregrine×red shaheen (F. peregrinus peregrinus×F. peregrinus babylonicus); 4=saker falcon (Falco cherrug); 5=gyr falcon (Falco rusticolus); 6=interspecific hybrid saker×gyr (F. cherrug×F. rusticolus)

was not significantly different between falcons from

different zones, suggesting that falcons in both areas were subjected to overall similar conditions. Moreover, these results indicate that the differences in climate and altitude across the study zones did not affect on membrane thickness.

Given the large variation in membrane thickness and its potential importance in the protection of the embryo, we encourage other researchers to include measurements on egg membranes in studies exploring eggshell thickness variation. We also suggest that researchers and field naturalists should be able to determine whether eggs are infertile, aborted or hatched based on collections of pieces of eggshells from falcon nests, which could avoid erroneous interpretations of nest failure due to factors not related to embryo development. We argue that an increased attention paid to membranes would be beneficial and suggest that field researchers should collect and store eggshells for further research when possible, since they are non-invasive indicators of environmental or microbial contamination, the nutrient source of females during egg formation, a source of DNA for genetic studies and an excellent source for the detection of viral infections (Pearce et al. 1997; Miller et al. 2003; Chao et al. 2007; Oppel et al. 2009).

Acknowledgements We would like to acknowledge the persons in charge and the workers of Roc Falcon especially Harald Kuespert, Stania Kuespert, Oscar Oliva Piferrer and Sandor Sebestyen for providing detailed information about the falcons, for helping to collect the eggs and measuring the birds. We thank Andy Gosler, Sven Thatje, Hugo Robles and one anonymous referee for constructive comments. We also thank to Benno Böer (UNESCO Office Doha, Qatar) for his support and encouragement to improve our knowledge about the biology and conservation of endangered falcons. This work was conducted on a contract of the Spanish National Research Council

(Agencia Estatal Consejo Superior de Investigaciones Científicas, CSIC; to AMC), the logistic support of the Ayuntamiento de Sanaüja and the Project MICIIN-CGL 2009-10652 (BOS).

References

- Ahlborn G, Sheldon BW (2005) Enzymatic and microbiological inhibitory activity in eggshell membranes as influenced by layer strains and age and storage variables. Poult Sci 84:1935– 1941
- Ar A, Rahn H, Paganelly CV (1979) The avian egg: mass and strength. Condor 81:331–337
- Bennett RS (1995) Relative sensitivity of several measures of eggshell quality to the stage of embryonic development. Bull Environ Contam Toxicol 54:428–431
- Blom J, Lilja C (2004) A comparative study of growth, skeletal development and eggshell composition in some species of birds. J Zool 262:361–369
- Bunck CM, Spann JV, Pattee OH, Fleming WJ (1985) Changes in eggshell thickness during incubation: implications for evaluating the impact of organochlorine contaminants on productivity. Bull Environ Contam Toxicol 35:173–182
- Burnham WA, Enderson JH, Boardman TJ (1984) Variation in peregrine falcon eggs. Auk 101:578–583
- Cade TJ, Lincer JL, White CM, Roseneau DG, Swartz LG (1971) DDE residues and eggshell changes in Alaskan falcons and hawks. Science 172:955–957
- Castilla AM, Herrel A, Díaz G, Francesca A (2007) Developmental stage affects eggshell breaking strength in two ground-nesting birds: the partridge (Alectoris rufa) and the quail (Coturnix japonica). J Exp Zool 307A:471–477
- Castilla AM, Martínez de Aragón J, Herrel A, Møller S (2009a) Eggshell thickness variation in Red-legged Partridge (Alectoris rufa) from Spain. Wilson J Ornithol 121:167–170
- Castilla AM, Herrel A, Van Dongen S, Furio N, Negro JJ (2009b) Determinants of egg shell strength in endangered raptors. J Exp Zool 311A:303–311
- Castilla AM, Herrel A, Robles H, Malone J, Negro JJ (2010) The effect of developmental stage on eggshell thickness variation in endangered falcon species. Zoology (in press)
- Chao MR, Hsien CH, Yeh CM, Chou SJ, Chu C, Su YC, Yu CY (2007) Assessing the prevalence of Salmonella enterica in poultry hatcheries by using hatched eggshell membranes. Poult Sci 86:1651–1655
- Cobb GP, Bargar TA, Pepper CB, Norman DM, Houlis PD, Anderson TA (2003) Using chorioallantoic membranes for non-lethal assessment of persistent organic pollutant exposure and effect in oviparous wildlife. Ecotoxicology 12:31–45
- Cramp S, Simmons KEL (1980) Handbook of the birds of Europe, the Middle East and North of Africa: the birds of the Western Palearctic Volume II. Oxford University Press, Oxford
- Ecay TW, Stewart JR, Blackburn DG (2004) Expression of calbindin-D-28K by yolk sac and chorioallantoic membranes of the corn snake, Elaphe guttata. J Exp Zool B Mol Dev Evol 302B:517–525
- Elaroussi MA, Deluca HF (1994) Calcium-uptake by chorioallantoic membrane effects of vitamins D and K. Am J Physiol Endocrinol Metab 30:837–841
- Falk K, Møller S (1990) Clutch size effects on eggshell thickness in the Peregrine Falcon and European Kestrel. Ornis Scand 21:265–269
- Falk K, Møller S, Mattox W (2006) A long-term increase in eggshell thickness of Greenlandic Peregrine Falcons Falco peregrinus tundrius. Sci Total Environ 355:127–134

- Feberwee A, de Wit JJ, Landman WJM (2009) Induction of eggshell apex abnormalities by Mycoplasma synoviae: field and experimental studies. Avian Pathol 38:77–85
- Felsenstein J (1985) Phylogenies and the comparative method. Am Nat 125:1–15
- Felsenstein J (2008) Comparative methods with sampling error and within-species variation: contrasts revisited and revised. Am Nat 171:713–725
- Fernie KJ, Shutt L, Letcher RJ, Ritchie IJ, Bird DM (2009) Environmentally relevant concentrations of DE-71 and HBCD alter eggshell thickness and reproductive success of American kestrels. Environ Sci Technol 43:2124–2130
- Gosler AG, Higham JP, Reynolds J (2005) Why are birds' eggs speckled? Ecol Lett 8:1105–1113
- Jagannath A, Shore RF, Walker LA, Ferns PN, Gosler AG (2008) Eggshell pigmentation indicates pesticide contamination. J Appl Ecol 45:133–140
- Kamata R, Takahashi S, Shimizu A, Shiraishi F (2006) Avian transgenerational reproductive toxicity test with in ovo exposure. Arch Toxicol 80:846–856
- Karlsson O, Clas Lilja C (2008) Eggshell structure, mode of development and growth rate in birds. Zoology 111:494–502. doi:10.1016/j.zool.2007.11.005
- Kutchai H, Steen JB (1971) Permeability of the shell and shell membranes of hens' eggs during development. Respir Phyiol 11:265–278
- Mao KM, Murakami A, Iwasawa A, Yoshizaki N (2007) The asymmetry of avian egg-shape: an adaptation for reproduction on dry land. J Anat 210:741–748
- Miller MM, Ealey KA, Oswald WB, Schat KA (2003) Detection of chicken anemia virus DNA in embryonal tissues and eggshell membranes. Avian Dis 47:662–671
- Oppel S, Powell AN, O'Brien DM (2009) Using eggshell membranes as a non-invasive tool to investigate the source of nutrients in avian eggs. J Ornithol 150:109–115
- Peakall DB, Lincer JL (1996) Do PCBs cause eggshell thinning? Environ Pollut 91:127–129
- Peakall DB, Lew TS, Springer AM, Walker WW, Riscbrough RW, Monk JG, Jarman WM, Walton BJ, Reynolds LM, Fyfe RW, Kiff LF (1983) Determination of DDE and PCB contents of peregrine falcon eggs: a comparison of whole egg measurements and estimates derived from eggshell membranes. Arch Environ Contam Toxicol 12:523–528
- Pearce JM, Fields RL, Scribner KT (1997) Nest materials as a source of genetic data for avian ecological studies. J Field Ornithol 68:471–481
- Pepper CB, Rainwater TR, Platt SG, Dever JA, Anderson TA, McMurry ST (2004) Organochlorine pesticides in chorioallantoic membranes of Morelet's crocodile eggs from Belize. J Wildl Dis 40:493–500
- Rahbek C (1993) Captive breeding—a useful tool in the preservation of biodiversity. Biodivers Conserv 2:426–437
- Rahn H, Ar A (1974) The avian egg: incubation time, water loss and nest humidity. Condor 76:147–152
- Romanoff AL, Romanoff AJ (1949) Biochemistry of the avian embryo. Wiley, New York

Solomon SE (1991) Egg and eggshell quality. Wolfe Publishing, Aylesbury

- Tuan RS, Ono T, Akins RE, Koide M (1991) Experimental studies on cultured, shell-less fowl embryos: calcium transport, skeletal development, and cardio-vascular functions. In: Deeming DC, Ferguson MWJ (eds) Egg incubation: its effect on embryonic development in birds and reptiles. Cambridge University Press, Cambridge, pp 419–433
- Tucker GM, Heath MF (1994) Birds in Europe: their conservation status, BirdLife Conservation Series 3. BirdLife International, Cambridge

Vanderstoep J, Richards JF (1970) The changes in eggshell strength during incubation. Poult Sci 49:276–285

- Wegner P, Kleinstäuber G, Baum F, Schilling F (2005) Long-term investigation of the degree of exposure of German peregrine falcons (Falco peregrinus) to damaging chemicals from the environment. J Ornithol 146:34–54
- Wilkin TA, Gosler AG, Garant D, Reynolds SJ, Sheldon BC (2009) Calcium effects on life-history traits in a wild population of the great tit (Parus major): analysis of long-term data at several spatial scales. Oecologia 159:463–472
- Yoshizaki N, Saito H (2002) Changes in shell membranes during development of quail embryos. Poult Sci 81:246–251