

Corrigendum

Corrigendum to “Melanin-based color variation in the Bearded Vulture suggests a thermoregulatory function”
[Comp. Biochem. Physiol. 149A (2008) 87–91]

Antoni Margalida ^{a,*}, Juan José Negro ^b, Ismael Galván ^c

^a *Bearded Vulture Study and Protection Group. Apdo. 43 E-25520 El Pont de Suert (Lleida) Spain*

^b *Estación Biológica de Doñana (CSIC). Department of Evolutionary Ecology. Avda. María Luisa s/n E-41013 Sevilla, Spain*

^c *Museo Nacional de Ciencias Naturales (CSIC). Department of Evolutionary Ecology. C/ José Gutiérrez Abascal, 2. E28006 Madrid, Spain*

Available online 13 February 2008

The authors regret that in the above article, errors were made in the sample size or percentages of the Results section (3.1.2. *Ear tufts* and 3.1.3. *Pectoral band*).

The correct sample size or percentages are as follows:

3.1.2. *Ear tufts*

Out of 67 *G. b. barbatus*, 77.6% showed well developed ear tufts, 22.4% little and none (0%) had it absent.

3.1.3. *Pectoral band*

G. b. meridionalis ($n=34$) showed a complete pectoral band (18%), intermediate (53%) or absent (29%).

DOI of original article: [10.1016/j.cbpa.2007.10.017](https://doi.org/10.1016/j.cbpa.2007.10.017).

* Corresponding author. Tel.: +34 973 690 386.

E-mail address: margalida@inf.entoron.es (A. Margalida).